



ACTIVE⁺

AB-CLO

ORAL DRENCH

Mineralised oral drench for the control and treatment of internal parasites in sheep. Including persistent activity against Barber's Pole (*Haemonchus Contortus*), and Liver Fluke (*Fasciola Hepatica*).



ACTIVE +2

Active Ingredients: 1g/L Abamectin, 50g/L Closantel (as sodium salt) with 0.5g/L Selenium & 2.2g/L Cobalt

**ABAMECTIN
CLOSANTEL**

USE ON:
SHEEP



TREATMENT & CONTROL OF:
INTERNAL PARASITES

AB-CLO

ORAL DRENCH

Active Ingredients: 1g/L Abamectin, 50g/L Closantel (as sodium salt) with 0.5g/L Selenium & 2.2g/L Cobalt.



BENEFITS

- Persistent activity for up to 42 days against Barber's Pole (*Haemonchus contortus*).
- Oral drench for the treatment and control of roundworms and liver fluke in sheep.
- Broad-spectrum – Oral Drench.
- Dual Active – Roundworm and Liver Fluke control.
- Developed in New Zealand for New Zealand.

ADMINISTRATION

By law the user must take due care, obtaining expert advice if necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Active+ Ab-Clo Oral Drench is ready to use through standard drenching equipment. Rinse gun with water immediately after use. A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by liveweight in each group (ewes, wethers, rams, lambs).

Do not underdose. Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

Correct drenching technique should be used. Check accuracy of drench guns regularly.

DOSE RATE: 1mL/5kg bodyweight delivering 0.2mg/kg Abamectin, 10mg/kg Closantel, 0.1mg/kg Selenium and 0.44mg/kg Cobalt.

DO NOT USE IN LAMBS UNDER 15kg OR 4 WEEKS OF AGE.

LIVEWEIGHT	DOSE VOLUME
15 kg	3mL
16 - 20kg	4mL
21 - 25kg	5mL
26 - 30kg	6mL
31 - 35kg	7mL
36 - 40kg	8mL
41 - 45kg	9mL
46 - 50kg	10mL
51 - 55kg	11mL
56 - 60kg	12mL
61 - 65kg	13mL
66 - 70kg	14mL
71 - 75kg	15mL
Over 75kg	add 1mL / 5 kg

WITHHOLDING PERIODS

MEAT: 56 days, MILK: 56 days.

EFFICACY

Active+ Ab-Clo Oral contains 1g/L Abamectin a broad spectrum drench of the macrocyclic lactone group, 50g/L Closantel (as sodium salt) a narrow spectrum salicylanilide drench and 0.5g/L Selenium and 2.2 g/L Cobalt. It has persistent activity up to 42 days against Barber's Pole (*Haemonchus contortus*).

At recommended doses, this product is effective against sensitive strains of the following mature and immature roundworms and liver fluke in sheep:

Barber's Pole (*Haemonchus contortus*), Stomach hair worm (*Trichostrongylus axei*), Black scour worm (*Trichostrongylus colubriformis* and *Trichostrongylus vitrinis*), Brown stomach worm (*Teladorsagia circumcincta* and *Teladorsagia trifurcata*), Small intestinal worm (*Cooperia oncophora*, *Cooperia surnabada* and *Cooperia punctata*), Thin-necked intestinal worm (*Nematodirus spathiger*), Liver fluke (*Fasciola hepatica*), Lungworm (*Dictyocaulus filaria*).

STORAGE

Store at or below 25°C. Protect from light. Stability in cold temperatures has not been assessed. Discard unused product.

PACK SIZES

Active+ Ab-Clo Oral is available in 20L, 10L and 5L packs.

Active+ Ab-Clo Oral is registered pursuant to the ACVM Act 1997 No. A011496.

09 274 8848 | www.nexan.co.nz

Registered to:
Nexan Corporation Limited. PO Box 217234. Botany Junction. Auckland

