



ACTIVE⁺

CYMOXI

POUR-ON DRENCH

Fast acting, long lasting, broad spectrum treatment & control of internal & external parasites of cattle (including lactating dairy cattle), and the treatment and control of Lungworm and Roundworm of Deer.



ACTIVE +1

MOXIDECTIN

Active Ingredient: 5mg/mL Moxidectin.

USE ON:
CATTLE
DEER



TREATMENT & CONTROL OF:
INTERNAL & EXTERNAL PARASITES,
LUNGWORM & ROUNDWORM

CYMOXI

POUR-ON DRENCH

Active Ingredient: 5mg/mL Moxidectin.



BENEFITS

- Contains Moxidectin - a potent macrocyclic lactone (ML).
- Improves milk production.
- Rainfast.
- Persistent activity.
- Non-Irritant formulation.
- Broad-spectrum.
- Safe to use concurrently with other treatments.
- Developed in New Zealand for New Zealand.

ADMINISTRATION

By law the user must take due care, obtaining expert advice if necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

SHAKE WELL BEFORE USE.

Invert drum three times before use. Administer 1mL per 10kg liveweight using any standard drenching equipment. Correct drenching technique should be used. Equipment should be checked for accuracy of dose delivery before drenching commences. Check accuracy of drench guns regularly.

DOSAGE RATE: 1mL per 10kg liveweight. This will administer the correct dose of 0.5mg Moxidectin/kg liveweight. The following table may be used as a guide for the dose volume to use.

| LIVEWEIGHT | DOSE VOLUME | DOSES PER PACK | | |
|------------|-------------|----------------|-----|------|
| | | 500 mL | 1L | 5L |
| 50 kg | 5 mL | 100 | 200 | 1000 |
| 60 kg | 6 mL | 83 | 166 | 833 |
| 70 kg | 7 mL | 71 | 142 | 714 |
| 80 kg | 8 mL | 62 | 125 | 625 |
| 90 kg | 9 mL | 55 | 111 | 555 |
| 100 kg | 10 mL | 50 | 100 | 500 |
| 110 kg | 11 mL | 45 | 90 | 454 |
| 120 kg | 12 mL | 41 | 83 | 416 |
| 130 kg | 13 mL | 38 | 76 | 384 |
| 140 kg | 14 mL | 35 | 71 | 357 |
| 150 kg | 15 mL | 33 | 66 | 333 |
| 200 kg | 20 mL | 25 | 50 | 250 |
| 250 kg | 25 mL | 20 | 40 | 200 |
| 300 kg | 30 mL | 16 | 33 | 166 |
| 350 kg | 35 mL | 14 | 28 | 142 |
| 400 kg | 40 mL | 12 | 25 | 125 |
| 450 kg | 45 mL | 11 | 22 | 111 |
| 500 kg | 50 mL | 10 | 20 | 100 |
| 550 kg | 55 mL | 9 | 18 | 90 |
| 600 kg | 60 mL | 8 | 16 | 83 |
| 650 kg | 65 mL | 7 | 15 | 76 |
| 700 kg | 70 mL | 7 | 14 | 71 |
| 750 kg | 75 mL | 6 | 13 | 66 |

For animals heavier than 750kg, use a volume of 1mL / 10kg liveweight.

WITHHOLDING PERIODS

MILK: NIL, MEAT: Cattle - NIL, Deer - NIL, Bobby Calf - NIL.

STORAGE

Store below 25°C. Protect from light.

RAIN

The efficacy of Active+ Cymoxi Pour-On is not adversely affected if applied when the hide is wet, or if rain occurs shortly after application. However treatment of animals under these circumstances is not recommended practice.

PERSISTANT ACTIVITY

Active+ Cymoxi Pour-On, following a single application at the recommended dose rate prevents reinfection with the following parasite species:

| CATTLE | | DEER | |
|--|------------------------------|----------------------------------|------------------------------|
| Parasite Species | Persistent protection period | Parasite Species | Persistent protection period |
| <i>Dictyocaulus viviparus</i> | 42 days | <i>Dictyocaulus ekerti</i> | 42 days |
| <i>Ostertagia ostertagi</i> (<i>O. lyrata</i>) | 35 days | <i>Ostertagia-like spp*</i> | 42 days |
| <i>Haemonchus spp</i> | 28 days | <i>Oesophagostomum venulosum</i> | 42 days |
| <i>Trichostrongylus axei</i> | 28 days | <i>Chabertia ovina</i> | 42 days |
| <i>Oesophagostomum radiatum</i> | 42 days | | |
| <i>Bunostomum phlebotomum</i> | 42 days | | |

**Ostertagia leptospicularis* and *Ostertagia-like species* including *Spiculoptera assymetrica*, *Spiculoptera spiculoptera* and *Skrjabinagia kolchida*.

PACK SIZES

Active+ Cymoxi Pour-On is available in 5L packs.

Active+ Cymoxi Pour-On is registered pursuant to the ACVM Act 1997 No. A011492.

09 274 8848 | www.nexan.co.nz

Registered to;
Nexan Corporation Limited. PO Box 217234. Botany Junction. Auckland

