



ACTIVE⁺

CYRO LIQUID

For the prevention of blowfly strike on sheep.



Active Ingredient: 500 g/L Cyromazine

USE ON:
SHEEP 

PREVENTION OF:
BLOWFLY STRIKE

CYRO LIQUID

LIQUID FOR SHEEP

Active Ingredient: 500 g/L Cyromazine



BENEFITS

- Water-soluble suspension for jetting, dipping and dressing
- Long term protection - up to 12 weeks
- Very competitive cost per dose
- Developed in New Zealand for New Zealand

ADMINISTRATION

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

USE RATE FOR LONGER TERM PROTECTION (UP TO 12 WEEKS)

Mix 2L of Active⁺ Cyro Liquid with 1000L of water and apply by showering, plunge dipping or jetting. Ensure saturation to skin level. Top up at the same rate. Because this is a non-stripping dip, no reinforcement is required.

USE RATE FOR MEDIUM TERM PROTECTION (UP TO 6 WEEKS)

Mix 1L of Active⁺ Cyro Liquid with 1000L of water and apply by showering or plunge dipping. Ensure saturation to skin level. Top up at the same rate. Because this is a nonstripping dip, no reinforcement is required. Do not use this rate for hand jetting or through spray races.

Refer to label for detailed dosing instructions.

WITHHOLDING PERIODS

MEAT: Sheep producing meat or offal for human consumption must not be sold for slaughter either during or within 7 days of the last treatment.

MILK: Milk intended for sale for human consumption must be discarded during treatment and for 35 days following the last treatment.

WOOL RESIDUES

Ensure the time between dipping and subsequent shearing is at least two months, except in the case of spot treatment of flystrike. For advice on your dipping programme, contact your supplier.

STORAGE

Store at or below 25°C, avoid temperatures below 0°C and above 35°C. Protect from light.

EFFICACY

Active⁺ Cyro Liquid is a non-stripping saturation dip which will give up to 12 weeks residual protection against flystrike caused by all commonly found strike flies including Australian blowfly (*Lucilia cuprina*).

A shorter period of protection may occur if Active⁺ Cyro Liquid is applied incorrectly, if there is excessive fleece soiling or if wool infections such as mycotic dermatitis (fleece rot) are present. Prolonged heavy rain following treatment may also reduce the protection period provided by Active⁺ Cyro Liquid.

PACK SIZES

Active⁺ Cyro Liquid is available in 10L packs.

Registered pursuant to the ACVM Act 1997, No. A011467.
See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to the HSNO Act 1996, HSR002233.
See www.epa.govt.nz for controls

09 274 8848 | www.nexan.co.nz

Registered to:
Nexan Corporation Limited, PO Box 217234, Botany Junction, Auckland