



ACTIVE⁺

EPI

POUR-ON FOR CATTLE AND DEER

For the treatment and control of internal and external parasites of beef and dairy cattle (including lactating cows) and internal parasites of deer.



ACTIVE +1

EPRINOMECTIN

Active Ingredient: 5g/L Eprinomectin.

USE ON:
CATTLE
DEER



TREATMENT & CONTROL OF:
INTERNAL & EXTERNAL PARASITES

EPI

POUR-ON FOR CATTLE AND DEER

Active Ingredient: 5g/L Eprinomectin.



BENEFITS

- For the treatment and control of internal parasites, lungworm and sucking lice in dairy and beef cattle.
- Potent Endectocide (Eprinomectin) - providing activity against a broad-spectrum of internal and external parasites.
- Zero milk, meat and bobby calf withholding periods and improves milk production.*
- Superior persistent activity against a broad-spectrum of parasites.
- Effective, safe and convenient.
- Weatherproof.
- Developed in New Zealand, for New Zealand.

*INCREASED MILK PRODUCTION - Increases Milk production in grazing dairy cows - Data has shown that milk production increases when cows or heifers infected with gastrointestinal nematode parasites are treated with an effective anthelmintic.

ADMINISTRATION

The dose is 1mL for each 10kg of liveweight to give 0.5mg Eprinomectin per kg liveweight. For pour-on application only. Apply along the topline in a narrow strip extending from the withers to the tailhead. Use the following dosage schedule. A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves etc).

Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing. Correct drenching technique should be used.

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

DOSE: 1mL for each 10kg of liveweight to give 0.5mg Eprinomectin per kg liveweight.

WEIGHT RANGE	DOSE VOLUME	DOSES / 5L PACK	WEIGHT RANGE	DOSE VOLUME	DOSES / 5L PACK
50 kg	5.0 mL	1000	251 - 300kg	30.0 mL	166
51 - 75kg	7.5 mL	666	301 - 350kg	35.0 mL	142
76 - 100kg	10.0 mL	500	351 - 400kg	40.0 mL	125
101 - 125kg	12.5 mL	400	401 - 450kg	45.0 mL	111
126 - 150kg	15.0 mL	333	451 - 500kg	50.0 mL	100
151 - 175kg	17.5 mL	285	501 - 550kg	55.0 mL	90
176 - 200kg	20.0 mL	250	551 - 600kg	60.0 mL	83
201 - 250kg	25.0 mL	200	601 - 650kg	65.0 mL	76

Heavier animals (for example, mature bulls) should receive an additional 5mL for each 50kg over 650kg. Consult your veterinarian or animal health adviser for assistance in the diagnosis, treatment and control of parasitism.

RAIN

Rainfall before or after treatment will not affect the efficacy of Active+ Epi Pour-On.

PACK SIZES

Active+ Epi Pour-On is available in 20L and 5L packs.

WITHHOLDING PERIODS

MILK: There is no milk withholding period for lactating dairy cattle. Milk from treated cows may be used for human consumption at any time following treatment with Active+ Epi Pour-On.

MEAT:

CATTLE - there is no meat withholding period for cattle.

DEER - producing meat and offal for human consumption must not be sold for slaughter either during treatment or within 7 days of the last treatment.

There is no withholding period for bobby calves born to cows treated with Active+ Epi Pour-On.

PERSISTENT ACTIVITY IN CATTLE

Active+ Epi Pour-On for cattle and deer when given at the recommended dosage of 1mL/10kg liveweight prevents reinfection of cattle with the following internal parasites for the time period indicated.

PARASITE	PERSISTENT ACTIVITY
<i>Ostertagia spp.</i>	28 days
<i>Cooperia spp.</i>	21 days
<i>Nematodirus helvetianus</i>	14 days
<i>Oesophagostomum radiatum</i>	28 days
<i>Dictyocaulus viviparus</i>	28 days
<i>Haemonchus contortus</i>	14 days
<i>Trichostrongylus spp.</i>	14 days

For best results Active+ Epi Pour-On for Cattle and Deer should be part of a total parasite control program. Check the effectiveness of anthelmintics at least annually. Follow recommended parasite management practices to reduce the potential for resistance to develop.

Consult your veterinarian for a complete parasite control program.

STORAGE

Store in the original container, tightly closed and locked up in a safe place. Store below 25°C away from direct sunlight.

Active+ Epi Pour-On is registered pursuant to the ACVM Act 1997 No. A011491.

09 274 8848 | www.nexan.co.nz

Registered to:
Nexan Corporation Limited. PO Box 217234. Botany Junction. Auckland

