



ACTIVE⁺

ABA

POUR-ON DRENCH

Fast acting, broad spectrum treatment & control of internal & external parasites of cattle.



ACTIVE +1

ABAMECTIN

Active Ingredient: 10 g/L Abamectin.

USE ON:
CATTLE 

TREATMENT & CONTROL OF:
INTERNAL & EXTERNAL PARASITES

ABA

POUR-ON DRENCH

Active Ingredient: 10 g/L Abamectin.



BENEFITS

- Improves milk production.*
- Convenient pour-on application.
- Broad-spectrum activity.
- Effective against sucking and biting lice (*Damalinea bovis*, *Linognathus vituli*).
- Effective against lungworm (adult and immaturely *Dictyocaulus viviparus*).
- Solvent base delivering high blood levels of Abamectin.
- Dramatic visible results.
- Developed in New Zealand for New Zealand.

*INCREASED MILK PRODUCTION - Increases milk production in grazing dairy cows - Data has shown that milk production increases when cows or heifers infected with gastrointestinal nematode parasites are treated with an effective anthelmintic.

ADMINISTRATION

Ready for use with standard pour-on equipment. Check accuracy of pour-on applicator regularly. A representative sample of animals should be weighed before treatment either with scales or weighband.

Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves etc). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

Apply along the backline from shoulders to tail.

This product should not be used on cattle that are under 100kg in weight or less than 16 weeks old.

CATTLE

DOSE: 1ml per 20kg bodyweight (500 micrograms Abamectin per kg).

WEIGHT RANGE	DOSE VOLUME
101 - 150 kg	7.5 mL
151 - 200 kg	10 mL
201- 250 kg	12.5 mL
251 - 300 kg	15 mL
301 - 350 kg	17.5 mL
351 - 400 kg	20 mL
401 - 450 kg	22.5 mL
451 - 500 kg	25 mL
Over 500 kg	1 mL/20 kg

RAIN

Treatment of animals with wet hides or when rain threatens should be avoided.

EFFICACY

Active+ Aba Pour-On for Cattle when used at the recommended dose level of 500 mg Abamectin per kg liveweight effectively controls the following parasites in cattle:

- Gastrointestinal roundworms: adult and immature *Ostertagia spp* (including inhibited immatures), *Haemonchus spp*, *Trichostrongylus spp*, *Cooperia spp*, *Bunostomum spp*, *Oesophagostomum spp*, and adults of *Chabertia spp*, *Nematodirus spp*, *Trichuris spp* and *Strongyloides spp*.
- Lungworms: adult and immature *Dictyocaulus viviparus*.
- Biting and Sucking Lice: *Bovicola bovis*, *Linognathus vituli*.

WITHHOLDING PERIODS

MEAT: Cattle producing meat and offal for human consumption must not be sold for slaughter during or within 35 days of the last treatment.

MILK: 35 days.

STORAGE

Store in the original container, tightly closed and locked up in a safe place. Store below 25°C away from direct sunlight.

PACK SIZES

Active+ Aba Pour-On is available in 20L and 5L packs.

Active+ Aba Pour-On is registered pursuant to the ACVM Act 1997 No. A011493.

09 274 8848 | www.nexan.co.nz

Registered to:
Nexan Corporation Limited, PO Box 217234, Botany Junction, Auckland

